

Solid-phase synthesis of terminal oligonucleotide-phosphoramidate conjugates

Cooke, L. A., Frauendorf, C., Gilea, M., Holmes, S. C., & Vyle, J. (2006). Solid-phase synthesis of terminal oligonucleotide-phosphoramidate conjugates. *Tetrahedron Letters*, 47(5), 719-722.
<https://doi.org/10.1016/j.tetlet.2005.11.098>

Published in:
Tetrahedron Letters

Document Version:
Peer reviewed version



Queen's University Belfast - Research Portal:
[Link to publication record in Queen's University Belfast Research Portal](#)


General rights

Copyright for the publications made accessible via the Queen's University Belfast Research Portal is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.



Take down policy

The Research Portal is Queen's institutional repository that provides access to Queen's research output. Every effort has been made to ensure that content in the Research Portal does not infringe any person's rights, or applicable UK laws. If you discover content in the Research Portal that you believe breaches copyright or violates any law, please contact openaccess@qub.ac.uk.




¹H nmr 1





^{31}P nmr 1


 ^{13}C nmr 1

Mass spectrum 1








**7b**




(N.B. Buffer A = 100 mM TEAA pH 6.5 - **NO** MeCN)







(**7b** after purification Buffer A = 5% MeCN, 100 mM TEAA pH 6.5)





**7d**



**7f**